

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Currently Amended) A process for ~~[stabilizing]~~ enhancing the solubility of a blood protein solution comprising:

(a) ~~[providing a blood protein solution;~~
(b)] adding to ~~[the]~~ a blood protein solution hydroxypropyl- α -cyclodextrin in an amount sufficient to form a stable complex with the protein; and

~~[(e)]~~ (b) lyophilizing the solution of step ~~[(b)]~~ (a) to form a lyophilized ~~[protein/hydroxypropyl- α -cyclodextrin]~~ complex of the protein and hydroxypropyl- α -cyclodextrin.

2. (Currently Amended) The process according to claim 1, further comprising reconstituting the lyophilized ~~[protein/hydroxypropyl- α -cyclodextrin]~~ complex.

3. (Currently Amended) The process according to claim 1, further comprising heating the blood protein solution, before or after adding hydroxypropyl- α -cyclodextrin, at a temperature of at least ~~[about]~~ 60°C for a time sufficient to inactivate any viruses present ~~[in the protein/hydroxypropyl- α -cyclodextrin]~~ complex.

4. (Currently Amended) The process according to claim 3 wherein the blood protein solution is heated for at least ~~[about]~~ 10 hours.

Appln No. 09/772,634

Amdt date August 11, 2003

Reply to Office action of February 11, 2003

5. (Currently Amended) The process according to claim 3 wherein the blood protein solution is heated to a temperature of at least ~~[about]~~ 80°C for at least ~~[about]~~ 72 hours.

6. (Currently Amended) The process according to claim 3 wherein the blood protein solution is heated to ~~[about]~~ at least 100°C for at least ~~[about]~~ 1 hour.

7. (Original) The process according to claim 1, further comprising subjecting the blood protein solution, before or after adding the hydroxypropyl- α -cyclodextrin, to a solvent detergent viral inactivation step.

8. (Original) The process according to claim 1, wherein the hydroxypropyl- α -cyclodextrin is present in the protein solution in an amount ranging from about 0.5% wt/vol. to about 15% wt/vol.

9. (Original) The process according to claim 1, wherein the hydroxypropyl- α -cyclodextrin is present in the protein solution in an amount ranging from about 1% wt/vol. to about 12% wt/vol.

10. (Currently Amended) The process according to claim 2, wherein the protein is present in the reconstituted ~~[protein/hydroxypropyl- α -cyclodextrin]~~ complex in an amount greater than about 0.1% wt/vol.

11. (Currently Amended) The process according to claim 2 wherein the protein is present in the reconstituted ~~[protein/hydroxypropyl- α -cyclodextrin]~~ complex in an amount from about 1% to about 8 %.

12. (Original) The process according to claim 1 wherein the protein is selected from the group consisting of albumin, Factor II, Factor VII, Factor VIII, Factor IX, Factors X and X_a, fibrinogen, antithrombin III, transferrin, haptoglobin, gamma globulins, fibronectin, protein C, protein S, thrombin and C1-inhibitor.

Appln No. 09/772,634

Amdt date August 11, 2003

Reply to Office action of February 11, 2003

13. (Original) The process according to claim 1, wherein the protein is fibrinogen.

14. (Original) The process according to claim 12, wherein the hydroxypropyl- α -cyclodextrin is present in the protein solution in an amount ranging from about 0.5% wt/vol. to about 15% wt/vol.

15. (Original) The process according to claim 12, wherein the hydroxypropyl- α -cyclodextrin is present in the protein solution in an amount ranging from about 2% wt/vol. to about 12% wt/vol.

16. (Currently Amended) The process according to claim 12, wherein the fibrinogen is present in the reconstituted [~~protein/hydroxypropyl- α -cyclodextrin~~] complex in an amount greater than about 1% wt/vol.

17. (Currently Amended) The process according to claim 12, wherein the protein is fibrinogen, and the fibrinogen is present in the reconstituted [~~protein/hydroxypropyl- α -cyclodextrin~~] complex in an amount from about 3% wt/vol. to about 10% wt/vol.

18. (Currently Amended) A process for [~~stabilizing~~] enhancing the stability of a fibrinogen solution comprising:

- (a) [~~providing a fibrinogen solution;~~
- (b)] adding to [~~the~~] a fibrinogen solution hydroxypropyl- α -cyclodextrin in an amount sufficient to form a stable complex with the protein;
- (c)] lyophilizing the solution of step [(b)] (a) to form a lyophilized [~~fibrinogen/hydroxypropyl- α -cyclodextrin~~] complex of fibrinogen and hydroxypropyl- α -cyclodextrin; and
- (d)] (c) reconstituting the lyophilized [~~fibrinogen/hydroxypropyl- α -cyclodextrin~~] complex.

Appln No. 09/772,634

Amdt date August 11, 2003

Reply to Office action of February 11, 2003

19. (Currently Amended) A lyophilized ~~[blood-protein/hydroxypropyl- α -cyclodextrin]~~ complex of a blood protein and hydroxypropyl- α -cyclodextrin prepared by:

(a) ~~[providing a blood protein solution;~~

~~(b)]~~ adding to ~~[the]~~ a blood protein solution hydroxypropyl- α -cyclodextrin in an amount sufficient to form a stable complex with the protein; and

~~[(e)]~~ (b) lyophilizing the solution of step ~~[(b)]~~ (a) to form the lyophilized ~~[blood-protein/hydroxypropyl- α -cyclodextrin]~~ complex.

20. (Currently Amended) A blood protein product prepared by:

(a) ~~[providing a blood protein solution;~~

~~(b)]~~ adding to ~~[the]~~ a blood protein solution hydroxypropyl- α -cyclodextrin in an amount sufficient to form a stable complex with the protein;

~~[(e)]~~ (b) lyophilizing the solution of step ~~[(b)]~~ (a) to form a lyophilized ~~[protein/hydroxypropyl- α -cyclodextrin]~~ complex of the protein and hydroxypropyl- α -cyclodextrin; and

~~[(d)]~~ (c) reconstituting the lyophilized ~~[protein/hydroxypropyl- α -cyclodextrin]~~ complex.

21. (Currently Amended) A fibrinogen product prepared by:

(a) ~~[providing a fibrinogen solution;~~

~~(b)]~~ adding to ~~[the]~~ a fibrinogen solution hydroxypropyl- α -cyclodextrin in an amount sufficient to form a stable complex with the protein;

~~[(e)]~~ (b) lyophilizing the solution of step ~~[(b)]~~ (a) to form a lyophilized ~~[fibrinogen/hydroxypropyl- α -cyclodextrin]~~ complex of fibrinogen and hydroxypropyl- α -cyclodextrin; and

~~[(d)]~~ (c) reconstituting the lyophilized ~~[fibrinogen/hydroxypropyl- α -cyclodextrin]~~ complex.

22. (Original) A blood protein product comprising a lyophilized solution of a stable complex of protein and hydroxypropyl- α -cyclodextrin.

23. (Currently Amended) The product according to claim 22, wherein the hydroxypropyl- α -cyclodextrin is present [~~in the solution~~] in an amount ranging from about 0.5% wt/vol. to about 15% wt/vol.

24. (Currently Amended) The product according to claim 22, wherein the hydroxypropyl- α -cyclodextrin is present [~~in the solution~~] in an amount ranging from about 1% wt/vol. to about 12% wt/vol.

25. (Original) The product according to claim 22, wherein the blood protein is fibrinogen.

26. (Original) A stabilized blood protein solution comprising a complex of the blood protein and hydroxypropyl- α -cyclodextrin.

27. (Original) The solution according to claim 26, wherein the protein is present in the complex in an amount greater than about 3% wt/vol.

28. (Currently Amended) The product according to claim 26, wherein the hydroxypropyl- α -cyclodextrin is present in the [~~solution~~] complex in an amount ranging from about 0.5% wt/vol. to about 15% wt/vol.

29. (Currently Amended) The process according to claim 26, wherein the hydroxypropyl- α -cyclodextrin is present in the [~~solution~~] complex in an amount ranging from about 1% wt/vol. to about 12% wt/vol.

30. (Original) The product according to claim 26, wherein the blood protein is fibrinogen.